



Investigation of vertex $a_1 \rightarrow VP$ for hadronic τ decays

K. Nurlan 

Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna, Russia

(E-mail: nurlan.qanat@gmail.com)

Abstract. In the framework of $U(3)$ symmetric quark model, triangular diagrams are calculated that describe the strongly interacting vertices $a_1 \rightarrow \rho\pi$ and $a_1 \rightarrow K^* K$. The divergences arising when considering quark loops are regularized by covariant cut off parameter. Based on four-particle hadronic τ decays, the quark structure of the a_1 resonance with quantum numbers $J^{PC}=1^{++}$ was studied. Theoretical estimates of intermediate channels with the a_1 meson for the partial decay widths of $\tau \rightarrow \pi\pi\pi\nu_\tau$, $\tau \rightarrow KK\pi\nu_\tau$ and $\tau \rightarrow KK\eta\nu_\tau$ are obtained. Contributions from contact channels describing the direct production of 3π , $KK\pi$ and $KK\eta$ mesons by τ lepton current are taken into account. The interference of contact and intermediate axial-vector channels in determining the integral decay widths is studied. The matrix elements of the processes are obtained in the leading approximation of $1/N_c$ expansion. It is shown that calculations on τ lepton mesonic decays confirm the quark-antiquark structure of the a_1 meson and clarify the role of the intermediate non-strange axial-vector channel in determining the integral partial decay widths. The obtained results are compared with experimental data by BaBar and Belle collaborations at the BEPC II and KEK lepton colliders.

Keywords: hadron structure, quark fields, meson interactions.

Introduction

Description of the mass spectrum and interactions of the axial-vector meson field a_1 is an important task in the non-strange $SU(2) \times SU(2)$ sector of low-energy QCD phenomenology. The properties and decays of the a_1 meson field were studied on the basis of inelastic reactions $\pi^+ p \rightarrow \pi^+ \pi^+ \pi^- p$ with a beam energy of 190 GeV/c by the COMPASS collaboration [1]. In the CLEO and LCHb experiments at the Large Hadron Collider, branching fractions of heavy $D^0 \rightarrow K^{\mp} \pi^{\pm} \pi^{\mp}$ meson decays measured, where the analysis of obtained data indicates the decisive role of intermediate channels with the a_1 meson [2, 3]. In addition, the parameters of the a_1 meson can be studied in the three-pion final state of the τ lepton decay. The latter comes without hadron impurity and, therefore, provides good accuracy. Relevant studies were carried out in [4,5].

The a_1 meson is characterized by quantum numbers $J^{PC}=1^{++}$. It is generally accepted that this meson is a bound state of $\bar{q}q$ quark fields. Quark fields are associated with the operator γ_{μ}, γ_5 . The mass spectrum of the a_1 meson in the ground and first excited states is described in the quark model [6].

In the present paper, we give a theoretical description of decays $a_1 \rightarrow \rho\pi$ and $a_1 \rightarrow K^* K$. The matrix elements of the indicated decays are calculated in the one-loop quark approximation in the leading order of expansion N_c , where N_c is the number of colors in QCD [6]. The estimates were obtained for the partial widths of $\tau \rightarrow \pi\pi\pi\nu_{\tau}$, $\tau \rightarrow KK\pi\nu_{\tau}$ and $\tau \rightarrow KK\eta\nu_{\tau}$ with different charge combinations which proceed predominantly through intermediate channels containing decays $a_1 \rightarrow \rho\pi$ and $a_1 \rightarrow K^* K$.

Methods

To describe the strong interaction vertices of meson a_1 with the $\rho\pi$ and $K^* K$ meson fields use the effective Lagrangian obtained on the basis of a model with four-quark interaction [6,7]. The decay $a_1 \rightarrow \rho\pi$ is described by the Feynman diagram shown in Fig. 1. The meson fields in the vertex $a_1 \rho\pi$ are coupled by the u and d quark fields. In the case of $a_1 K^* K$, the strange quark field is in the internal lines. The Lagrangian determined the interaction of meson fields with quarks is represented in the following form

$$\Delta L_{int} = \bar{q} [1/2 \gamma^{\mu} \gamma^5 \lambda_{\pm,0}^{a_1} g_{\rho} a_{1\mu}^{\pm,0} + i \gamma^5 \lambda_{\pm,0}^{\pi} g_{\pi} \pi^{\pm,0} (1) + 1/2 \gamma^{\mu} \lambda_{\pm,0}^{\rho} g_{\rho} \rho_{\mu}^{\pm,0} + 1/2 \gamma^{\mu} \lambda^K g_{K^*} K^*_{\mu} + i \gamma^5 \lambda_{\pm,0}^K g_K K^{\pm,0}] q,$$

where a_1, π, ρ, K^* and K are the meson fields; γ_5, γ_{μ} are Dirac matrices in four-dimensional Minkowski space; q and \bar{q} are u, d and s quark triplets with constituent masses $m_u \approx m_d = 270 \text{ MeV}$ and $m_s = 420 \text{ MeV}$ [7]; λ are linear combinations of Gell-Mann matrices

$$\lambda_{\pm}^K = \frac{\lambda_4 - i \lambda_5}{\sqrt{2}}, \quad \lambda_0^K = \frac{\lambda_6 + i \lambda_7}{\sqrt{2}},$$

$$\lambda_{\pm}^{\rho} = \frac{\lambda_1 - i \lambda_2}{\sqrt{2}}, \quad \lambda_0^{\rho} = \frac{\sqrt{2} \lambda_3 - \lambda_8}{\sqrt{2}}.$$




Figure 1. Triangle quark diagram describing decay $a_1 \rightarrow \rho\pi$

The coupling constants g_ρ , g_K , g_π and g_{K^*} determine the interactions of mesons with quarks [7]. These constants determine the renormalization of meson fields and are expressed through logarithmically divergent integrals

$$g_{K^*} = \sqrt{3/2 I_{11}}, \quad g_K = \sqrt{Z_K/4 I_{11}}, \quad (2)$$

$$g_\rho = \sqrt{3/2 I_{20}}, \quad g_\pi = \sqrt{Z_\pi/4 I_{20}},$$

where

$$I_{n_1 n_2} = -i \frac{N_C}{(2\pi)^4} \int \frac{\Theta(\Lambda_4^2 + k^2)}{(m_u^2 - k^2)^{n_1} (m_s^2 - k^2)^{n_2}} d^4 k. \quad (3)$$

Using the Lagrangian (1) above, one can construct a vertex with a triangular loop. Consideration of this vertex leads to an integral of the following form

$$I_{21}^{a_1 \rho \pi} = N_C \int \frac{\text{Tr}[\gamma^\mu \gamma^5 (\hat{k} + \hat{p}_\rho + m_s) \gamma^\nu (\hat{k} + m_u) \gamma^5 (\hat{k} - \hat{p}_\pi + m_u)] d^4 k}{[(k + p_\rho)^2 - m_u^2][k^2 - m_u^2][(k + p_\pi)^2 - m_u^2]} \frac{d^4 k}{(2\pi)^4}. \quad (4)$$

When calculating this integral, we use one-loop approximation [6]. As a result, obtain the following decay matrix element of $a_1 \rightarrow \rho\pi$

$$M(a_1 \rho \pi) = -i F_\pi g_\rho^2 Z_\pi \varepsilon_\mu(a_1) g_{\mu\nu} \varepsilon_\nu(\rho), \quad (5)$$

where $\varepsilon_\mu(a_1)$ and $\varepsilon_\nu(\rho)$ are polarization vectors.

In a similar way, we obtain the amplitude describing the decay $a_1 \rightarrow K^* K$ with the participation of strange mesons

$$M(a_1 K^* K) = -i \frac{3g_\rho g_K}{\sqrt{2}g_{K^*}} (3m_u - m_s) \varepsilon_\mu(a_1) g_{\mu\nu} \varepsilon_\nu(K^*). \quad (6)$$

Results and discussions

The contributions of axial-vector channels to hadronic τ decays

In this chapter we consider mesonic τ decays $\tau \rightarrow \pi\pi\pi\nu_\tau$, $\tau \rightarrow KK\pi\nu_\tau$ and $\tau \rightarrow KK\eta\nu_\tau$, where the vertices $a_1 \rightarrow \rho\pi$ and $a_1 \rightarrow K^*\pi$ obtained above contribute as intermediate channels. The diagrams corresponding to these channels are shown in Fig. 2 and 3. These channels are considered in conjunction with the contact diagram with the axial-vector part of the W boson, when the final products are produced directly by the lepton current. The corresponding diagram for the contact channel is shown in Fig. 4. The decay of $\tau \rightarrow W\nu_\tau$ is described by the electroweak Lagrangian of the standard model (Weinberg – Salam theory) [8,9]. The parts of diagrams describing W boson interaction with meson fields are obtained within the effective quark model.




Figure 2. Feynman diagram describing channel with a_1 meson for the decay $\tau \rightarrow 3\pi\nu_\tau$




Figure 3. Feynman diagram describing channel with a_1 meson for the decay $\tau \rightarrow KK\pi\nu_\tau$

The decay amplitude $\tau \rightarrow \pi\pi\pi\nu_\tau$ can be represented as the product of the lepton current with the hadronic current in the following form:

$$M(\tau \rightarrow \pi^+\pi^-\pi^-\nu_\tau) = -iF_\pi G_F V_{ud} L_\mu M_{A\mu} \quad (7)$$

Where $L_\mu = \bar{\nu}_\tau \gamma_\mu (1 - \gamma_5) \tau$ is the lepton current, G_F is the Fermi constant, V_{ud} is the element of the Cabibbo-Kobayashi-Maskawamatrix.

The total contribution from contact term and intermediate axial-vector channel reads

$$M_{A\mu} = Z_\pi g_\rho^2 (g_{\mu\nu} + ((q^2 - 6m_u^2)g_{\mu\nu} - q_\mu q_\nu) BW_{a_1}) BW_\rho (p_{\pi^+} - p_{\pi^-}^{(1)})_\nu + (p_{\pi^-}^{(1)} \leftrightarrow p_{\pi^-}^{(2)}), \quad (8)$$

where p_{π^+} , $p_{\pi^-}^{(1)}$ and $p_{\pi^-}^{(2)}$ are the momentum of charged pions in the final state.




Figure 4. The contact diagram describing decay $\tau \rightarrow KK\pi\nu_\tau$

The transition of the W boson into intermediate a_1 meson has a gauge-invariant structure

$$g^{\mu\nu} [q^2 - 6m_u^2] - q_\mu q_\nu \quad (9)$$

The a_1 meson in the intermediate state is described by the Breit-Wigner propagator

$$BW_M = \frac{g_{\mu\nu}}{M_{a_1}^2 - q^2 - i\sqrt{q^2}\Gamma_{a_1}}. \quad (10)$$

Table 1. Calculated branching fractions (Br) for hadronic τ decays. Comparison of theoretical results with experimental data of BaBar and Belle collaborations is presented.

Mode	$Br(\times 10^{-3})$				
	$\tau \rightarrow \pi^- \pi^+ \pi^- \nu_\tau$	$\tau \rightarrow \pi^0 \pi^0 \pi^- \nu_\tau$	$\tau \rightarrow K^- K^0 \pi^0 \nu_\tau$	$\tau \rightarrow K^- K^+ \pi^- \nu_\tau$	$\tau \rightarrow K^- K^0 \eta \nu_\tau$
Theor	90.5	91.5	1.46	0.4	0.0014
PDG [10]	90.2 ± 0.5	92.6 ± 1.0	1.50 ± 0.07	1.43 ± 0.02	< 0.009
BaBar	88.3 ± 1.3 [11]	-	-	1.34 ± 0.03 [11]	-
Belle	84.2 ± 2.5 [4]	-	1.49 ± 0.07 [12]	1.55 ± 0.01 [4]	-

To calculate the decays $\tau \rightarrow KK\pi\nu_\tau$ and $\tau \rightarrow KK\eta\nu_\tau$ consider the vertices of W boson interaction with meson fields presented Fig.3. When considering these vertices, use approximation in one-loop. The full matrix elements for axial-vector channels with an intermediate a_1 meson can be found in paper [13].

The obtained results for the decay channels $\tau \rightarrow 3\pi\nu_\tau$, $\tau \rightarrow KK\pi\nu_\tau$ and $\tau \rightarrow KK\eta\nu_\tau$ are given in Table 1. The uncertainty of theoretical calculations is estimated at 15% [6,7].

Conclusion

In the present paper, the vertices $a_1 \rightarrow \rho\pi$ and $a_1 \rightarrow K^* K$ going through the triangular quark loop are calculated (Fig.1). The amplitudes of the processes considered are obtained using chiral Lagrangians in leading order of $1/N_c$, where N_c is the number of colors in QCD. Estimates are obtained for the branching fractions of $\tau \rightarrow 3\pi\nu_\tau$, $\tau \rightarrow KK\pi\nu_\tau$ and $\tau \rightarrow KK\eta\nu_\tau$ taking into account the contact channel and channels with the intermediate meson a_1 . Interference between contact and axial vector channels is always negative due to the negative value of the Breit-Wigner propagator for most of the energy range $(\sum M_p)_2 \leq q^2 \leq M\tau^2$.

Our calculations for the considered hadronic τ decays confirm the quark-antiquark structure of the a_1 meson and clarify the role of intermediate channels with the nonstrange a_1 meson in determining the partial widths of the decays. Model predictions for the branching fraction do not contradict the experimental data of BaBar [11] and Belle [4,12].

Acknowledgement

The author thanks collaborators prof. M.K. Volkov and A.A. Pivovarov. This work was supported by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan Grant no. BR20280986.

The contribution of the authors

Kanat Nurlan – writing the text and critically revising its content, approval of the final version of the article for publication, agreement to be responsible for all aspects of the work, proper examination and resolution of issues related to the reliability of the data.

References

- Aghasyan M. et al. [COMPASS], Light isovector resonances in $\pi\text{-p} \gg \pi\text{-}\pi\text{-}\pi\text{-p}$ at 190 GeV/c // Phys. Rev. D., Vol. 98, (2018) No.9, 092003, doi:10.1103/PhysRevD.98.092003
- d'Argent P., Skidmore N., Benton J. et al. Amplitude Analyses of $D0 \gg \pi\text{-}\pi\text{-}\pi\text{-}\pi\text{-}$ and $D0 \gg K\text{-}K\text{-}\pi\text{-}\pi\text{-}$ Decays // JHEP, Vol. 05, (2017) No. 143, doi:10.1007/JHEP05(2017)143
- Aaij R. et al. [LHCb], Studies of the resonance structure in $D0 \gg K\pi\pi$ decays, Eur. Phys. J. C, Vol. 78, (2018) no.6, pp. 443, doi:10.1140/epjc/s10052-018-5758-4
- Lee M.J. et al. [Belle] // Measurement of the branching fractions and the invariant mass distributions for $\tau \gg h\text{-}h\text{-}h\text{-}\nu$ decays, Phys. Rev. D, Vol. 81, (2010) 113007, doi:10.1103/PhysRevD.81.113007
- Abreu P. et al. [DELPHI], A Measurement of the tau topological branching ratios // Eur. Phys. J. C. Vol. 20 (2001), pp. 617-637, doi:10.1007/PL00011152
- Ebert D., Reinhardt H., and Volkov M.K., Effective hadron theory of QCD // Prog. Part. Nucl. Phys., Vol. 33 (1994), pp. 1-120
- Volkov M.K. and Radzhabov A.E. The Nambu-Jona-Lasinio model and its development // Phys. Usp., Vol. 49 (2006), pp. 551-561, doi:10.1070/PU2006v049n06ABEH005905

Weinberg S., A Model of Leptons // Phys. Rev. Lett., Vol. 19 (1967), pp. 1264-1266, doi:10.1103/PhysRevLett.19.1264

Salam A. Weak and Electromagnetic Interactions // Conf. Proc. C. Vol. 680519 (1968), pp. 367-377, doi:10.1142/9789812795915_0034

Workman R.L. et al. [Particle Data Group], Review of Particle Physics // PTEP 2022 (2022), pp. 083C01, doi:10.1093/ptep/ptac097

Aubert B. et al. [BaBar], Exclusive branching fraction measurements of semileptonic tau decays into three charged hadrons, $\tau \rightarrow \phi \pi \nu(\tau)$ and $\tau \rightarrow \phi K \nu(\tau)$ // Phys. Rev. Lett., Vol. 100 (2008), pp. 011801, doi:10.1103/PhysRevLett.100.011801

Ryu S. et al. [Belle], Measurements of Branching Fractions of τ Lepton Decays with one or more $K0$ // Phys. Rev. D, Vol. 89, (2014) no.7, pp. 072009, doi:10.1103/PhysRevD.89.072009

Volkov M.K., Pivovarov A.A. and Nurlan K., Three-meson τ decays involving kaons and η mesons in the NJL model // Phys. Rev. D, Vol. 109, (2024) no.1, pp. 016016, doi:10.1103/PhysRevD.109.016016

Нурлан Қанат

*Н.Н. Боголюбов атындағы теориялық физика лабораториясы,
Біріккен ядролық зерттеулер институты, Дубна, Ресей
(E-mail: nurlan.qanat@gmail.com)*

τ лептонның адронды ыдыраулары үшін $a_1 \rightarrow VP$ шыңын зерттеу

Аңдатпа. $U(3)$ симметриялы кваркті модел шеңберінде пәрменді әсерлесу жүзеге асатын $a_1 \rightarrow \rho \pi$ және $a_1 \rightarrow K^* K$ шыңдарын сипаттайтын үшбұрышты диаграммалар есептелді. Кваркты диаграммаларды қарастырған кезде пайда болатын жинақсыздық ковариантты кесу параметрі арқылы реттелген. τ – лептонның төрт бөлшекті адрондық ыдырауы негізінде $J^{PC}=1^{++}$ кванттық сандары бар a_1 резонансының кварк құрылымы зерттелді. a_1 мезоны бар аралық арналардың теориялық бағалары $\tau \rightarrow \pi \pi \pi \nu$, $\tau \rightarrow K K \pi \nu$ және $\tau \rightarrow K K \eta \nu$ жартылай ыдырау ендері үшін есептелді. τ – лептонды токтан 3π , $K K \pi$ және $K K \eta$ мезондарының тікелей тууларын сипаттайтын байланыс арналарының үлестері есепке алынды. Ыдыраулардың интегралдық ендерін анықтауда контактілі және аралық аксиал-векторлы арналардың интерференциялары зерттелді. Процестердің матрицалық элементтері $1/N_c$ кеңеюінің жетекші жуықтауында алынды. τ – лептонның мезондық ыдыраулары бойынша есептеулер a_1 мезонының кварк-антикварк құрылымын растайтыны және ыдыраулардың ішінара интегралдық ендерін анықтаудағы аксиал-векторлық аралық арнаның рөлін нақтылайтыны көрсетілген. Алынған нәтижелер BEPC II және KEK лептон коллайдерлерінде алынған BaBar және Belle тәжірибелік деректерімен салыстырылды.

Түйін сөздер: адрондардың құрылымы, кваркты өрістер, мезондардың әсерлесулері.

Нурлан Канат

Лаборатория теоретической физики им. Н.Н. Боголюбова,
Объединенный институт ядерных исследований, Дубна, Россия
(E-mail: nurlan.qanat@gmail.com)

Исследование вершины $a_1 \rightarrow VP$ для адронных τ распадов

Аннотация. В рамках $U(3)$ симметричной кварковой модели вычислены треугольные диаграммы, описывающие сильно взаимодействующие вершины $a_1 \rightarrow \rho\pi$ и $a_1 \rightarrow K^*K$. Расходимости, возникающие при рассмотрении кварковых петель регуляризуются ковариантным обрезанием. На основе четырех частичных адронных распадов тау-лептона исследована кварковая структура резонанса a_1 с квантовыми числами $J^{PC}=1^{++}$. Получены теоретические оценки промежуточных каналов с мезоном a_1 для парциальных ширин распадов $\tau \rightarrow \pi\pi\pi\nu_\tau$, $\tau \rightarrow KK\pi\nu_\tau$ и $\tau \rightarrow KK\eta\nu_\tau$. Учтены вклады от контактных каналов, описывающие прямое рождение мезонов 3π , $KK\pi$ и $KK\eta$ тау-лептонным током. Исследованы интерференций контактных и промежуточных аксиально-векторных каналов в определении интегральных ширин распадов. Матричные элементы процессов получены в лидирующем приближении разложения $1/N_c$. Показано, что расчеты по мезонным распадам тау-лептона подтверждают кварк-антикварковую структуру мезона a_1 и проясняет роль промежуточного нестранного аксиально-векторного канала в определении парциальных интегральных ширин распадов. Полученные результаты сравниваются с экспериментальными данными BaBar и Belle, полученные на лептонных коллайдерах VEPP-II и KEK.

Ключевые слова: структуры адронов, кварковые поля, взаимодействия мезонов.

Information about the authors:

Нурлан Канат – PhD, Н.Н. Боголюбов атындағы теориялық физика лабораториясы, Біріккен ядролық зерттеулер институты, Дубна, Ресей.

Nurlan Kanat – PhD, researcher at Bogoliubovlaboratory of theoretical physics, Joint institute for nuclear research, Dubna, Russia

Нурлан Канат – PhD, научный сотрудник Лаборатории теоретической физики им. Н.Н. Боголюбова, Объединенный институт ядерных исследований, Дубна, Россия.



Copyright: © 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>).